

Small codimension smooth subvarieties in even-dimensional homogeneous spaces with Picard group $\mathbb{Z}$

Nicolas Perrin^a

^a*Université Pierre et Marie Curie - Paris 6
Institut de Mathématiques de Jussieu
175 rue du Chevaleret, 75013 Paris.*

Abstract

We investigate a method proposed by E. Arrondo and J. Caravantes to study the Picard group of a smooth low-codimension subvariety X in a variety Y when Y is homogeneous. We prove that this method is strongly related to the signature σ_Y of the Poincaré pairing on the middle cohomology of Y . We give under some topological assumptions a bound on the rank of Picard group $\text{Pic}(X)$ in terms of σ_Y and remove these assumptions for grassmannians to recover the main result of E. Arrondo and J. Caravantes.

1. Introduction

E. Arrondo and J. Caravantes present in [1] a sketch of program to decide if a divisor D in a smooth subvariety X of a smooth even-dimensional variety Y is equivalent to a multiple of an hyperplane section of X . To apply this method, they need some hard topological results that we will not discuss here. We will however discuss the general validity of the rest of the method in the case of homogeneous varieties Y with Picard group $\mathbb{Z}$ and prove that it is closely related to the signature σ_Y of the Poincaré pairing on the middle cohomology of Y . Let us fix the following notations: $N = \dim(Y)$ the dimension of Y is even and to keep coherent notations with [1] we set $N = 2(n - 1)$. Recall the definition (see [5]):

Definition 1.1 *A subvariety X of Y is said to be cumbersome if for any subvariety $Z \subset Y$ such that $\dim(Z) \geq \text{codim}(X)$ we have $[X] \cdot [Z] \neq 0$.*

We will prove the following:

Theorem 1.2 *Let X be a cumbersome smooth subvariety in Y of dimension n' with $n' \geq n$. Assume that X is simply connected and that, for any n -dimensional Schubert subvariety Y' of Y , the intersection of X with a general translate of Y' is irreducible, then $\text{Pic}(X)$ is free of rank at most $\frac{1}{2}(h^{n-1}(Y, \mathbb{C}) - \sigma_Y) + 1$.*

One major difficulty to use this result is to prove the two topological assumptions on X , namely that X is simply connected and that its intersection with the general translate of a Schubert subvariety of dimension n is irreducible. However, in the case where Y is the Grassmannian variety $\mathbb{G}(p, m)$, O. Debarre [5] proved such results. In that case, set $R = \max(p, m - p)$, $r = \min(p, m - p)$, $\delta = R - 1$ for $r \geq 3$ and $\delta = 1$ for $r \leq 2$. We deduce the following:

Email address: nperrin@math.jussieu.fr (Nicolas Perrin).

Corollary 1.3 *Assume X is a smooth cumbersome subvariety of dimension $n' \geq n - 1 + \delta$ of Y then $\text{Pic}(X)$ is free of rank at most $\frac{1}{2}(h^{n-1}(Y, \mathbb{C}) - \sigma_Y) + 1$.*

Finally we recover the main result in [1]:

Corollary 1.4 *Assume that $r \leq 2$, or equivalently that Y is a projective space or a Grassmannians of lines, then any smooth cumbersome subvariety X of Y of dimension $n' \geq n$ satisfies $\text{Pic}(X) = \mathbb{Z}$.*

Remark that stronger results were proved by A. J. Sommese [10] but in smaller codimension. In this note we prove that the method of E. Arrondo and J. Caravantes to show that the Picard group of X is $\mathbb{Z}$ will not work in general but only if the Poincaré duality is positive definite on the middle cohomology. The only¹ even-dimensional homogeneous varieties with Picard group $\mathbb{Z}$ and positive definite Poincaré pairing on the middle cohomology are the projective spaces, the grassmannians of lines, the quadrics of dimension multiple by 4 and the Cayley plane $\mathbb{O}\mathbb{P}^2$. It is striking that these are examples of projective spaces over composition algebras (over $\mathbb{C}$, $\mathbb{H}$ and $\mathbb{O}$, cf. [4]). With this in mind we ask the following:

Question 1.1 Is it true that, for any smooth cumbersome subvariety X of dimension at least 9 of the Cayley plane $\mathbb{O}\mathbb{P}^2$, we have $\text{Pic}(X) = \mathbb{Z}$?

Acknowledgements: I would like to thank L. Gruson for many usefull discussion during the preparation of this text.

2. Proof of the Theorem

To prove the theorem we will in some sense restrict ourselves to the n -dimensional case. Let X' be a variety satisfying the hypothesis of Theorem 1.2. Let D' be a divisor of X' , we may assume D' to be smooth². We will denote by X a general hyperplane section of X' of dimension n and by D the divisor in X corresponding to D' . Remark that X and D are also smooth and that X is cumbersome.

It is a classical result that a basis of $H^*(Y, \mathbb{Z})$ is given by the classes $\sigma(w)$ of Schubert subvarieties where w is in a coset W_Y of the Weyl group W (see for example [3]). We will denote by $(\sigma(t))_{t \in T}$ and $(\sigma(u))_{u \in U}$ the classes of Schubert subvarieties of dimensions n and $n - 1$. They form a basis of the effective monoids in $H^{n-2}(Y, \mathbb{Z})$ and $H^{n-1}(Y, \mathbb{Z})$. We use the notation: $\sigma(w) \cdot \sigma(w') = \sum_{w''} c_{w, w'}^{w''} \sigma(w'')$ and write

$$[X] = \sum_{t \in T} a_t \sigma(t) \quad \text{and} \quad [D] = \sum_{u \in U} \alpha_u \sigma(u)$$

with a_t and α_u non negative for all $t \in T$ and $u \in U$ (because X is cumbersome, all the a_t are positive).

Poincaré duality acts as a permutation between Schubert classes and induces an involution $w \mapsto w^*$ on W_Y . In particular, it stabilises U and we may consider the partition $U_{id} \coprod U_{hyp} = U$ where U_{id} is the set of fixed points of U under the Poincaré involution. This induces a decomposition of the abelian group $H^{n-1}(Y, \mathbb{Z})$ into $H_{id}^{n-1}(Y, \mathbb{Z}) \oplus H_{hyp}^{n-1}(Y, \mathbb{Z})$. Poincaré duality acts as the identity on $H_{id}^{n-1}(Y, \mathbb{Z})$ and is hyperbolic on $H_{hyp}^{n-1}(Y, \mathbb{Z})$. In particular if σ_Y is its signature on $H^{n-1}(Y, \mathbb{Z})$, we have the equalities $\dim(H_{id}^{n-1}(Y, \mathbb{Z})) = \sigma_Y$ and $\dim(H_{hyp}^{n-1}(Y, \mathbb{Z})) = h^{n-1}(Y, \mathbb{Z}) - \sigma_Y$.

We first compare D and H_X with respect to numerical equivalence. Consider the classes of curves $C(u) = \sigma(u) \cdot [X]$ and $C(t) = [D] \cdot \sigma(t)$. We define some non negative integers by: $x_u := H_X \cdot C(u)$, $y_u := D \cdot C(u)$, $z_t := H_X \cdot C(t)$ and $\lambda_t := [D] \cdot C(t)$. We have the equalities:

¹ See [9]

² This will be harmless by replacing D' by $D' + mH_{X'}$ with m big.

$$x_u = \sum_{t \in T} c_{t,H}^{u*} a_t, \quad y_u = \alpha_{u*} \quad \text{and} \quad z_t = \sum_{u \in U} c_{t,H}^{u*} \alpha_u.$$

Define a matrix M with two lines and columns indexed by $U \cup T$ by

$$M = (m_{i,j})_{i \in \{1,2\}, j \in U \cup T} = \begin{pmatrix} (x_u)_{u \in U} & (z_t)_{t \in T} \\ (y_u)_{u \in U} & (\lambda_t)_{t \in T} \end{pmatrix}.$$

The same proof as in [1] and Lefschetz's hyperplane Theorem lead to the following³:

Proposition 2.1 *The divisor D is numerically equivalent to a multiple of H_X iff M is of rank one.*

PROOF — We follow the proof of E. Arrondo and J. Caravantes in [1]. If D is numerically equivalent to a multiple of H_X , then the rank of the matrix M is one. Conversely, if the second line of the matrix is q times the first one, let $D_0 = [D] - qH_X$. The intersections $D_0 \cdot C(u)$ and $D_0 \cdot C(t)$ vanish. But H_X^{n-1} and H_X^{n-2} are linear combinations of $C(u)$ and $C(t)$ respectively, hence if S is a smooth surface obtained from X by $n-2$ hyperplane sections, we get that

$$D_0|_S H|_S = D_0 H_X^{n-1} = 0 \quad \text{and} \quad D_0|_S^2 = D_0(D - qH_X)H_X^{n-2} = 0.$$

By Hodge index Theorem, the divisor $D_0|_S$ has to be numerically trivial. By Lefschetz's hyperplane Theorem, this has to be true for D_0 . $\square$

Corollary 2.2 *The divisor D' is numerically equivalent to a multiple of $H_{X'}$ iff M is of rank one.*

PROOF — We know from the previous Proposition that D and H_X are numerically dependent. Lefschetz's hyperplane theorem implies that it is also the case for D' and $H_{X'}$. $\square$

As in [1], the sequence $0 \rightarrow N_{D/X} \rightarrow N_{D/Y} \rightarrow (N_{X/Y})|_D \rightarrow 0$ is exact because X and D are smooth. Taking the top Chern classes gives the equality $P := D \cdot_Y D - (D \cdot_X D) \cdot_X X|_X = 0$. In terms of the variables $(a_t)_{t \in T}$, $(\alpha_u)_{u \in U}$ and $(\lambda_t)_{t \in T, v \in V}$, we get:

$$P = \sum_{u \in U} \alpha_u \alpha_{u*} - \sum_{t \in T} a_t \lambda_t.$$

The next step is the elimination of the variables $(\lambda_t)_{t \in T, v \in V}$. We consider for this, following [1], the surfaces $(S_t)_{t \in T}$ where S_t is the intersection of X with a general subvariety of class $\sigma(t)$. We set:

$$\text{Hodge}_t := \begin{vmatrix} H_{S_t}^2 & H_{S_t} D_{S_t} \\ H_{S_t} D_{S_t} & D_{S_t}^2 \end{vmatrix} = \begin{vmatrix} \sum_{u \in U} c_{t,H}^u x_u & z_t \\ \sum_{u \in U} c_{t,H}^u y_u & \lambda_t \end{vmatrix} = \begin{vmatrix} \sum_{u \in U} c_{t,H}^u x_u & \sum_{u \in U} c_{t,H}^u y_u \\ \sum_{u \in U} c_{t,H}^u y_u & \lambda_t \end{vmatrix} = \sum_{u \in U} c_{t,H}^u \begin{vmatrix} m_{1,u} & m_{1,t} \\ m_{2,u} & m_{2,t} \end{vmatrix}.$$

Observe that Hodge_t is a non negative linear combination of some minors of the matrix M . By hypothesis on X' , the surface S_t is irreducible hence Hodge_t is non positive. Set $d_t := H_{S_t}^2 > 0$ and define the following quadratic form q in the variables $(\alpha_u)_{u \in U}$ (observe that q is non positive)

$$q := P + \sum_{t \in T} \frac{a_t}{d_t} \text{Hodge}_t. \tag{1}$$

In [1], E. Arrondo and J. Caravantes decompose q , in the special case of grassmannians of lines, as a sum of squares to prove its non negativity. We do this in full generality and show that some negative part

³ We can apply a numerical version of Lefschetz Theorem because X' and all its linear sections are simply connected thanks to the homotopic Lefschetz's hyperplane Theorem, see [7, Theorem 3.1.21]

may appear. We know that all the variables $(x_u)_{u \in U}$ do not vanish because X is cumbersome. Hence we may define two quadratic forms q' and q'' in the variables $(\alpha_u)_{u \in U}$ by:

$$q' = \frac{1}{2} \sum_{t \in T} \sum_{(u, u') \in U^2} \frac{a_t c_{t,H}^u c_{t,H}^{u'}}{d_t x_u x_{u'}} \begin{vmatrix} m_{1,u} & m_{1,u'} \\ m_{2,u} & m_{2,u'} \end{vmatrix}^2 \quad \text{and} \quad q'' = \frac{1}{2} \sum_{u \in U} \frac{1}{x_u x_{u^*}} \begin{vmatrix} m_{1,u} & m_{1,u^*} \\ m_{2,u} & m_{2,u^*} \end{vmatrix}^2.$$

Proposition 2.3 *The formula $q = q' - q''$ holds.*

PROOF — For any couple (u, u') of elements in U , set $A_{u,u'} = - \sum_{t \in T} \frac{a_t}{d_t} c_{t,H}^{u^*} c_{t,H}^{u'^*}$.

The coefficient of $\alpha_u \alpha_{u'}$ in q is equal to $A_{u,u'} + A_{u',u} = 2A_{u,u'}$ if $u' \neq u^*$ and $u' \neq u$, it is equal to $2 + 2A_{u,u'}$ if $u' = u^* \neq u$, to $A_{u,u'}$ if $u^* \neq u' = u$ and to $1 + A_{u,u'}$ if $u' = u^* = u$. The coefficient of $\alpha_u \alpha_{u'}$ in q' is equal to $2A_{u,u'}$ if $u' \neq u$ and, if $u' = u$, to

$$\sum_{t \in T} \sum_{u'' \in U, u'' \neq u} \frac{a_t}{d_t} c_{t,H}^{u^*} c_{t,H}^{u''^*} \frac{x_{u''^*}}{x_{u^*}} = \sum_{t \in T} \frac{a_t}{d_t x_{u^*}} c_{t,H}^{u^*} (d_t - c_{t,H}^{u^*} x_{u^*}) = \frac{x_u}{x_{u^*}} - \sum_{t \in T} \frac{a_t}{d_t} c_{t,H}^{u^*}^2 = \frac{x_u}{x_{u^*}} + A_{u,u'}.$$

The non vanishing coefficients of $\alpha_u \alpha_{u'}$ in q'' are equal to -2 if $u' = u^* \neq u$ and to $\frac{x_u}{x_{u^*}}$ if $u' = u \neq u^*$. But this is exactly the difference $q' - q''$. $\square$

Corollary 2.4 *If for all $u \in U$, we have $\begin{vmatrix} m_{1,u} & m_{1,u^*} \\ m_{2,u} & m_{2,u^*} \end{vmatrix} = 0$, then D' is a multiple of $H_{X'}$ in $\text{Pic}(X')$.*

PROOF — Indeed, in that case q'' vanishes so q must vanish (and also q'). Because X is cumbersome, all a_t are different from 0 and this implies that all 2×2 minors of M vanish and M is of rank one. Apply Corollary 2.2 to conclude that D' is numerically equivalent to a multiple of H . Then because X' is simply connected, this implies the result in the Picard group of X' . $\square$

We finish the proof of Theorem 1.2. Because X' is simply connected, we know that the Picard group is free of finite type. Let us now prove that its rank is at most $\frac{1}{2}(h^{n-1}(Y) - \sigma_Y) + 1$. Indeed, let $(D_i)_{0 \leq i \leq k}$ some divisors on X' independant in $\text{Pic}(X')$ with $D_0 = H$. Let us consider the conditions in Corollary 2.4. These conditions are $x_u \alpha_u - x_{u^*} \alpha_{u^*} = 0$ for $u \in U$ and in fact there are only $\frac{1}{2}h_{hyp}^{n-1}(Y) = \frac{1}{2}(h^{n-1}(Y) - \sigma_Y)$ such conditions because this condition is trivial if $u = u^*$ and is the same for u and u^* . Now if k is strictly bigger than the preceding number of conditions, then there exists an element D' in the linear span of the family $(D_i)_{1 \leq i \leq k}$ satisfying these conditions. But then D' as to be a multiple of $H = D_0$, a contradiction.

3. Applications

Denote by $Y = \mathbb{G}(p, m)$ the Grassmannian variety of p -dimensional vector subspaces in a m -dimensional vector space. Thanks to [5, Corollaire 7.4] and [5, Théorème 8.1 and Corollaire 8.3], we obtain ⁴

Proposition 3.1 (i) *Let X be a cumbersome irreducible subvariety in $\mathbb{G}(p, m)$ of dimension n' such that $2n' \geq N + r$ (or equivalently $n' \geq n - 1 + \frac{r}{2}$) then X is simply connected.*

(ii) *Let X be a cumbersome irreducible variety of dimension $n' \geq n - 1 + \delta$ in Y , then the intersection of X with a general translate of a Schubert variety of dimension n is irreducible.*

⁴ The first part of this result generalises the celebrated Fulton-Hansen Theorem [6] while the second generalises Bertini's Theorem for Grassmannians.

These propositions complete the proof of Corollaries 1.3 and 1.4⁵

We may not hope that the method proposed by E. Arrondo and J. Caravantes will lead to a better bound on the rank of the Picard group because the decomposition into sums of squares shows that some negative terms appear. More precisely we have the following

Proposition 3.2 *If Y is cominuscule, for any square of minor appearing q'' the coefficient in q' of this square is smaller than its coefficient in q'' .*

PROOF — If t is such that $c_{t,H}^u c_{t,H}^{u*} \neq 0$ then the quiver of t (see [8] for more on these quivers) differs from those of u and u^* by one vertex and has to be the union of these quivers. In particular t is unique. Furthermore we get $d_t \geq x_u + x_{u^*} \geq 2a_t$ thus $\frac{a_t}{d_t} \leq \frac{1}{2} < 1$. $\square$

For example, consider the case $Y = Q_{2(n-1)}$ is a smooth quadric. Then our result together with Fulton-Hansen connectivity Theorem and Bertini Theorem leads to the:

Proposition 3.3 *If X is a smooth subvariety of dimension $n' \geq n$ in a smooth quadric $Q_{2(n-1)}$ of dimension $2(n-1)$, then $\text{Pic}(X) = \mathbb{Z}$ if n is odd and $\text{Pic}(X) = \mathbb{Z}$ or $\mathbb{Z}^2$ if n is even.*

This has already been observed by E. Arrondo and J. Caravantes, and in the last case, they give in [1] an example of smooth subvariety X of dimension n in $Q_{2(n-1)}$ with Picard group $\mathbb{Z}^2$.

References

- [1] Arrondo E., Caravantes J., *On the Picard group of low-codimension subvarieties*, arXiv:math.AG/0511267.
- [2] Barth W., Larsen M. E., *On the homotopy groups of complex projective algebraic manifolds*. Math. Scand. 30 (1972), 88–94.
- [3] Brion M., *Lectures on the geometry of flag varieties*. Topics in cohomological studies of algebraic varieties, 33–85, Trends Math., Birkhäuser, Basel, 2005.
- [4] Chaput P.E., *Geometry over composition algebras: projective geometry*, J. Algebra 298 (2006), no. 2, 340–362.
- [5] Debarre O., *Théorèmes de connexité pour les produits d'espaces projectifs et les grassmanniennes*. Amer. J. Math. 118 (1996), no. 6, 1347–1367.
- [6] Fulton W., Hansen J., *A connectedness theorem for projective varieties, with applications to intersections and singularities of mappings*. Ann. of Math. (2) 110 (1979), no. 1, 159–166.
- [7] Lazarsfeld, R. *Positivity in algebraic geometry. I*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, 48. Springer-Verlag, Berlin, 2004. xviii+387 pp.
- [8] Perrin N., *Small resolutions of minuscule Schubert varieties*, math.AG/0601117 to appear in Compositio Mathematica.
- [9] Slodowy, P., *On the signature of homogeneous spaces*. Geom. Dedicata 43 (1992), no. 1, 109–120.
- [10] Sommese A. J., *Complex subspaces of homogeneous complex manifolds. II. Homotopy results*. Nagoya Math. J. 86 (1982), 101–129.

⁵ The signature σ_Y is easy to compute in the last case, see [9].